



# Texas Medicaid/CHIP Vendor Drug Program

## Drug Utilization Criteria For Outpatient Use Guidelines

### Platelet Aggregation Inhibitors

#### About

Information on indications for use or diagnosis is assumed to be unavailable. All criteria may be applied retrospectively; prospective application is indicated with an asterisk [\*]. The information contained is for the convenience of the public. The Texas Health and Human Services Commission is not responsible for any errors in transmission or any errors or omissions in the document.

#### Publication History

Developed December 2016.

---

### 1. Dosage [\*]

#### *Adults*

Platelet aggregation inhibitors (PAIs) are FDA-approved to reduce thrombotic cardiovascular events in patients with a history of ischemic stroke, or to prevent stroke in patients with predisposing factors for atherosclerosis or symptomatic cerebrovascular disease.<sup>1-11</sup> PAIs work by interfering with pathways that promote normal platelet function: inhibiting cyclooxygenase-1 (e.g., aspirin); inhibiting adenosine uptake into platelets, resulting in increased cyclic-3',5'-adenosine monophosphate (cAMP) and adenosine levels (e.g., dipyridamole); inhibiting the adenosine diphosphate (ADP) P2Y<sub>12</sub> receptor on the platelet surface and blocking activation of the glycoprotein GPIIb/IIIa complex (e.g., clopidogrel, prasugrel, ticagrelor); antagonizing protease-activated receptor 1 (PAR-1), which inhibits thrombin and thrombin receptor agonist peptide activity (e.g., vorapaxar); or inhibiting phosphodiesterase III (e.g., cilostazol).<sup>2-4, 12</sup>

Aspirin is available in combination with omeprazole, a proton pump inhibitor, to reduce the risk of aspirin-associated gastric ulcers in those patients requiring aspirin for secondary prevention of cardiovascular and cerebrovascular events.<sup>2-4, 10</sup> Aspirin is also available as combination therapy with dipyridamole, pairing two antiplatelet agents with different mechanisms of action for secondary stroke prevention.<sup>2-4, 11</sup> Maximum recommended adult dosages for PAIs are summarized in Table 1. Medication profiles identifying patients prescribed dosages exceeding these recommendations will be reviewed.



**Texas Medicaid/CHIP Vendor Drug Program**  
**Drug Utilization Criteria For Outpatient Use Guidelines**  
**Platelet Aggregation Inhibitors**

**Table 1: Maximum Daily Adult Dosages for PAIs<sup>1-3, 5-11</sup>**

<b>Drug</b>	<b>Dosage Form/Strength</b>	<b>Maximum Recommended Daily Dose</b>
<i>Monotherapy</i>		
aspirin (Durlaza®)	162.5 mg extended-release capsule	<i>reduce risk of death and recurrent stroke or recurrent MI in patients with a history of ischemic stroke or TIA, and/or a history of chronic coronary artery disease: 162.5 mg once daily</i>
cilostazol (generics)	50 mg, 100 mg tablets	<i>intermittent claudication: 100 mg twice daily</i>
clopidogrel (Plavix®, generics)	75 mg, 300 mg tablets	<i>acute coronary syndrome (NSTEMI-ACS and STEMI): following a 300 mg loading dose, 75 mg/day in combination with aspirin</i>  <i>thromboembolism prophylaxis in patients with recent MI or stroke, or established peripheral vascular disease: 75 mg/day</i>
dipyridamole (generics)	25 mg, 50 mg, 75 mg tablets	<i>prevention of postoperative thrombotic complications in patients with prosthetic heart valves: 400 mg/day (divided doses, in combination with warfarin) or 300 mg/day (divided doses, in combination with aspirin)</i>
prasugrel (Effient®)	5 mg, 10 mg tablets	<i>acute coronary syndrome in patients to be managed with PCI: following a 60 mg loading dose, 10 mg/day<sup>+</sup> in combination with aspirin</i>
ticagrelor (Brilinta®)	60 mg, 90 mg tablets	<i>acute coronary syndrome: following a 180 mg loading dose, 90 mg twice daily<sup>^</sup> in combination with aspirin or clopidogrel</i>
vorapaxar (Zontivity®)	2.08 mg tablet	<i>MI, stroke, thrombosis prophylaxis in patients with a history of MI or peripheral arterial disease: 2.08 mg/day in combination with aspirin or clopidogrel</i>
<i>Combination Therapy</i>		
aspirin/omeprazole (Yosprala®)	81 mg/40 mg, 325 mg/40 mg delayed-release tablets	<i>secondary prevention of cardiovascular and cerebrovascular events in patients predisposed to gastric ulcers: 325 mg/40 mg once daily</i>
dipyridamole/aspirin (Aggrenox®, generics)	200 mg/25 mg extended-release capsule	<i>stroke prevention: 200 mg/25 mg twice daily</i>

MI = myocardial infarction; TIA = transient ischemic attack; NSTEMI-ACS = non-ST-segment elevation acute coronary syndrome; STEMI = ST-segment elevation myocardial infarction; PCI = percutaneous coronary intervention

<sup>+</sup>patients < 60 kg may use prasugrel 5 mg/day as maintenance therapy in combination with aspirin to reduce bleeding risk

<sup>^</sup>ticagrelor dosages are decreased to 60 mg twice daily after 12 months

**Pediatrics**

Dipyridamole is FDA-approved for use in pediatric patients 12 years of age and older as adjunctive therapy to prevent thromboembolism following cardiac valve replacement. The maximum recommended dose is 100 mg four times daily in combination with warfarin. Dosages exceeding these recommendations will be reviewed.



# Texas Medicaid/CHIP Vendor Drug Program

## Drug Utilization Criteria For Outpatient Use Guidelines

### Platelet Aggregation Inhibitors

Aspirin as Durlaza®, cilostazol, prasugrel, ticagrelor, vorapaxar, aspirin/omeprazole, and dipyridamole/aspirin are not recommended for use in pediatric patients as safety and efficacy have not been established for these agents in this patient population. Although not FDA-approved, clopidogrel has effectively been used in pediatric patients to reduce thrombosis risk in infants and children with select types of heart disease, or as an alternative in patients with Kawasaki disease or ischemic stroke when aspirin is not tolerated.<sup>2, 3, 13-16</sup>

## 2. Duration of Therapy

There is no basis for limiting PAI therapy duration when prescribed to prevent thromboembolic events associated with cardiovascular or cerebrovascular disease. However, PAI therapy duration varies, based on medication utilized and indication for use. PAI treatment durations are summarized in Table 2.

<b>Table 2: Maximum Treatment Duration for PAIs in Adults<sup>1-3, 5-11, 17, 18</sup></b>		
<b>Drug</b>	<b>Indication</b>	<b>Maximum Treatment Duration</b>
<i>Monotherapy</i>		
aspirin (Durlaza®)	<i>reduce risk of death and recurrent stroke or recurrent MI in patients with a history of ischemic stroke or TIA, and/or a history of chronic coronary artery disease</i>	indefinite
cilostazol (generics)	<i>intermittent claudication</i>	indefinite
clopidogrel (Plavix®, generics)	<i>acute coronary syndrome (NSTEMI-ACS and STEMI)</i>	up to 1 year, in combination with aspirin; aspirin then continued indefinitely <sup>+</sup>
	<i>thromboembolism prophylaxis</i>	indefinite
dipyridamole (generics)	<i>prevention of postoperative thrombotic complications in patients with prosthetic heart valves</i>	indefinite, in combination with warfarin or aspirin
prasugrel (Effient®)	<i>acute coronary syndrome in patients to be managed with PCI</i>	at least 12 months, in combination with aspirin, after stent placement
ticagrelor (Brilinta®)	<i>acute coronary syndrome</i>	90 mg twice daily x 1 year in combination with aspirin; then, 60 mg twice daily in combination with aspirin indefinitely
vorapaxar (Zontivity®)	<i>MI, stroke, thrombosis prophylaxis in patients with a history of MI or peripheral arterial disease</i>	indefinite, in combination with aspirin or clopidogrel
<i>Combination Therapy</i>		
aspirin/omeprazole (Yosprala®)	<i>secondary prevention of cardiovascular and cerebrovascular events in patients predisposed to gastric ulcers</i>	indefinite
dipyridamole/aspirin (Aggrenox®, generics)	<i>stroke prevention</i>	indefinite

MI = myocardial infarction; TIA = transient ischemic attack; NSTEMI-ACS = non-ST-segment elevation acute coronary syndrome; STEMI = ST-segment elevation myocardial infarction; PCI = percutaneous coronary intervention

<sup>+</sup>in patients with aspirin allergy, clopidogrel monotherapy may be continued indefinitely



# Texas Medicaid/CHIP Vendor Drug Program

## Drug Utilization Criteria For Outpatient Use Guidelines

### Platelet Aggregation Inhibitors

### 3. Duplicative Therapy [\*]

Combined administration of multiple PAIs may result in augmented antiplatelet effects. Adjunctive therapy with aspirin and clopidogrel, dipyridamole, prasugrel, ticagrelor, or vorapaxar has documented efficacy for acute coronary syndrome or thrombotic event prevention; concurrent therapy with clopidogrel and ticagrelor or vorapaxar is also FDA-approved for thromboembolic event prophylaxis or acute coronary syndrome (see Tables 1 and 2). Patients receiving concurrent therapy, however, should be observed for enhanced adverse effects due to similar mechanisms of actions between PAIs.<sup>1-11, 17, 18</sup>

### 4. Drug-Drug Interactions [\*]

Patient profiles will be assessed to identify those drug regimens which may result in clinically significant drug-drug interactions. Major drug-drug interactions considered clinically significant for platelet aggregation inhibitors are summarized in Table 3. Only those drug-drug interactions classified as clinical significance level 1/contraindicated or those considered life-threatening which have not yet been classified will be reviewed.

**Table 3: Major PAI Drug-Drug Interactions<sup>1-3, 5-11, 19-21</sup>**

Target Drug	Interacting Drug	Interaction	Recommendation	Clinical Significance*
PAIs, including aspirin	low molecular weight heparins	potential for additive bleeding adverse effects; PAIs inhibit platelet aggregation and have increased bleeding risk, prolonged bleeding time	avoid concurrent therapy, if possible; if drug combination necessary, use cautiously, monitor for signs/symptoms of bleeding	major, moderate (DrugReax) 2-major, 3-moderate (CP)
PAIs, including aspirin	selective serotonin reuptake inhibitors (SSRIs)/ serotonin norepinephrine reuptake inhibitors (SNRIs)	increased bleeding risk with combined therapy; SSRIs/SNRIs deplete platelet serotonin, which may impair platelet aggregation	monitor for signs/symptoms of bleeding; may consider substituting tricyclic antidepressant for SSRI/SNRI	SSRIs –major; SNRIs-major (DrugReax) 3-moderate (CP)
PAIs, including aspirin	warfarin	combined administration may increase bleeding risk, due to additive effects	if combined therapy necessary, monitor patients closely for bleeding signs/symptoms	major (DrugReax) 2-major, 3-moderate (CP)
aspirin	methotrexate (MTX)	potential for increased MTX serum levels, risk of enhanced pharmacologic/toxic effects as NSAIDs can reduce MTX clearance	avoid concurrent NSAIDs within 10 days of high-dose MTX; otherwise, use cautiously together; monitor for increased myelosuppressive, GI adverse effects; may consider using longer leucovorin rescue	major (DrugReax) 1-severe (CP)



**Texas Medicaid/CHIP Vendor Drug Program**  
**Drug Utilization Criteria For Outpatient Use Guidelines**  
**Platelet Aggregation Inhibitors**

**Table 3: Major PAI Drug-Drug Interactions (continued)**<sup>1-3, 5-11, 19-21</sup>

Target Drug	Interacting Drug	Interaction	Recommendation	Clinical Significance*
clopidogrel	dasabuvir/ ombitasvir/ paritaprevir/ ritonavir (Viekira®)	adjunctive administration with clopidogrel (strong CYP2C8 inhibitor) contraindicated by manufacturer, as dasabuvir is metabolized by CYP2C8, which increases risk for dasabuvir-induced QT interval prolongation; ritonavir, a CYP3A4 inhibitor, may limit clopidogrel conversion to active metabolite	avoid concurrent use	1-severe (CP)
cilostazol, dipyridamole	riociguat (Adempas®)	concurrent administration may result in increased hypotension risk	avoid concurrent use	contraindicated (DrugReax) dipyridamole: 1-severe; cilostazol: 3-moderate (CP)
cilostazol, ticagrelor, vorapaxar	itraconazole, strong CYP3A4 inhibitors	co-administration may result in elevated serum concentrations of select PAIs and potential bleeding complications as cilostazol, ticagrelor, and vorapaxar metabolized by CYP3A4	avoid use; ticagrelor therapy should not be initiated for at least 2 weeks after itraconazole discontinuation; if adjunctive administration necessary, use cautiously and monitor patient closely for enhanced pharmacologic/adverse effects, especially bleeding	ticagrelor: contraindicated; cilostazol, vorapaxar: major (DrugReax) ticagrelor: 1-severe; cilostazol, vorapaxar: 2-major (CP)

\*CP = Clinical Pharmacology; PAI – platelet aggregation inhibitor

**REFERENCES**

1. DRUGDEX® System (electronic version). Truven Health Analytics, Greenwood Village, Colorado, USA. Available at: <http://www.micromedexsolutions.com.libproxy.uthscsa.edu>. Accessed December 8<sup>th</sup>, 2016.
2. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2016. Available at: <http://www.clinicalpharmacology.com>. Accessed December 8<sup>th</sup>, 2016.
3. Facts and Comparisons eAnswers [database online]. Hudson, Ohio: Wolters Kluwer Clinical Drug Information, Inc.; 2016. Available at: <http://eanswers.factsandcomparisons.com.ezproxy.lib.utexas.edu/>. Accessed December 8<sup>th</sup>, 2016.
4. Cucchiara BL, Messé SR. Antiplatelet therapy for secondary prevention of stroke. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. (Accessed on December 8<sup>th</sup>, 2016.)
5. Aspirin extended-release capsules (Durlaza®) package insert. New Haven Pharmaceuticals, September 2015.



## Texas Medicaid/CHIP Vendor Drug Program

### Drug Utilization Criteria For Outpatient Use Guidelines

#### Platelet Aggregation Inhibitors

6. Clopidogrel tablets (Plavix®) package insert. Sanofi-Aventis, July 2015.
7. Prasugrel tablets (Effient®) package insert. Eli Lilly and Company, July 2016.
8. Ticagrelor (Brilinta®) package insert. AstraZeneca, September 2016.
9. Vorapaxar tablets (Zontivity®) package insert. Merck and Co., Inc., April 2015.
10. Aspirin/omeprazole extended-release tablets (Yosprala®) package insert. Aralez Pharmaceuticals US Inc., September 2016.
11. Aspirin/extended-release dipyridamole capsules (Aggrenox®) package insert. Boehringer Ingelheim Pharmaceuticals, Inc., November 2015.
12. Coutre S. Congenital and acquired disorders of platelet function. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. (Accessed on December 8<sup>th</sup>, 2016.)
13. Li JS, Yow E, Berezny KY, et al. Dosing of clopidogrel for platelet inhibition in infants and young children: primary results of the Platelet Inhibition in Children On cLOpidogrel (PICOLO) trial. *Circulation* 2008;117:553-9.
14. Soman T, Rafay MF, Hune S, et al. The risks and safety of clopidogrel in pediatric arterial ischemic stroke. *Stroke*. 2006;37(4):1120-2.
15. Giglia TM, Massicotte MP, Tweddell JS, et al., for the American Heart Association Congenital Heart Defects Committee of the Council on Cardiovascular Disease in the Young, Council on Cardiovascular and Stroke Nursing, Council on Epidemiology and Prevention, and Stroke Council. Prevention and treatment of thrombosis in pediatric and congenital heart disease: a scientific statement from the American Heart Association. *Circulation*. 2013;128:2622-2703.
16. Newburger JW, Takahashi M, Gerber MA, et al. Diagnosis, treatment, and long-term management of Kawasaki disease: a statement for health professionals from the Committee on Rheumatic Fever, Endocarditis and Kawasaki Disease, Council on Cardiovascular Disease in the Young, American Heart Association. *Circulation*. 2004;110:2747-71.
17. Spinler SA, deDenus S. Chapter 7. Acute coronary syndromes. (Chapter) In: DiPiro JT, Talbert RL, Yee GC, et al. (eds): *Pharmacotherapy: a pathophysiologic approach*. 9<sup>th</sup> edition. New York: McGraw-Hill; 2014. Access Pharmacy Website. Available at: <http://accesspharmacy.mhmedical.com.ezproxy.lib.utexas.edu/book.aspx?bookid=689>. Accessed December 9, 2016.
18. Hoebe BJ, Talbert RL. Chapter 12. Peripheral arterial disease. (Chapter) In: DiPiro JT, Talbert RL, Yee GC, et al. (eds): *Pharmacotherapy: a pathophysiologic approach*. 9<sup>th</sup> edition. New York: McGraw-Hill; 2014. Access Pharmacy Website. Available at: <http://accesspharmacy.mhmedical.com.ezproxy.lib.utexas.edu/book.aspx?bookid=689>. Accessed December 9, 2016.
19. Ombitasvir, paritaprevir, and ritonavir tablets; dasabuvir tablets co-packaged for oral use (Viekira Pak®) package insert. AbbVie Inc., June 2016.
20. Dasabuvir, ombitasvir, paritaprevir, and ritonavir extended-release tablets (Viekira XR) package insert. AbbVie Inc., July 2016.
21. DRUG-REAX® System (electronic version). Thomson Reuters (Healthcare) Inc., Greenwood Village, Colorado, USA. Available at: <http://www.thomsonhc.com.libproxy.uthscsa.edu>. **Accessed December 9<sup>th</sup>, 2016.**

#### Prepared by

- Drug Information Service, the University of Texas Health Science Center at San Antonio.
- The College of Pharmacy, the University of Texas at Austin.